

Special Reprint

WORLD JOURNAL OF  
**ORTHODONTICS**


Volume 11 • Number 1 • 2010

**EFFICIENCY AND EFFECTIVENESS  
OF SURESMILE**

*Alana K. Saxel/Lenore J. Louie/James Mah*

**suressmile**

OFFICIAL JOURNAL OF THE



WORLD FEDERATION OF ORTHODONTISTS

# EFFICIENCY AND EFFECTIVENESS OF SURESMILE

Alana K. Saxe, DMD<sup>1</sup>

Lenore J. Louie, MSc,  
DMD<sup>2</sup>

James Mah, DDS, MSc,  
DMSc<sup>3</sup>

**Aim:** To examine the efficiency and effectiveness of the SureSmile process using the standards of the American Board of Orthodontists Objective Grading System (ABO OGS). **Methods:** Three diplomates of the American Board of Orthodontics provided study casts of 62 patients whose orthodontic treatment was consecutively completed. Patients treated using the SureSmile process and a conventional approach were anonymized and randomized prior to independent scoring by two ABO OGS-calibrated examiners. **Results:** Intra- and interexaminer reliability was consistent in all components with no differences between examiners ( $r = 0.96$ ,  $P < .001$ ). The ABO OGS score for the SureSmile patients (mean 26.3) was 4.4 points lower ( $P < .001$ ) than for those treated conventionally (mean 30.7). Furthermore, treatment with the SureSmile process was shorter (14.7 months vs 20.0 months). **Conclusion:** The SureSmile process results in a lower mean ABO OGS score and a reduced treatment time than conventional approaches. The approach has great potential to both decrease treatment time and improve quality. *World J Orthod* 2010;11:16–22.

**Key words:** computer, efficiency, orthodontics, treatment

Efficient management of orthodontic patients in delivering timely care is an important aspect of treatment.<sup>1</sup> Orthodontics is focused on developing methods or techniques to decrease treatment time while maintaining quality outcomes. Although orthodontists have pursued this goal for quite some time, at present, there is no conventional orthodontic biomechanical treatment approach that has demonstrated any greater efficiency or effectiveness than any other. This is likely due to the complexity of orthodontics and the diversity of patients. Thus, it is imperative to understand the factors that can adversely impact the length of orthodontic treatment and use this information to implement measures that promote care more predictably and timely.

Studies have shown that average treatment time ranges from 23.1 to 31.2 months, depending on the design of the investigation and other factors.<sup>1,2</sup> A study from New Zealand determined average treatment length to be 23.5 months with a range of 12.0 to 37.0

months (SD 4.7,  $n = 366$ ),<sup>3</sup> similar to the previously reported 23.1 months<sup>2</sup> and 22.0<sup>4</sup> months from a comparable study. However, longer treatment times (28.6 months<sup>5</sup> and 31.2 months<sup>1</sup>) are also found in the relevant literature.

Patient cooperation is well-recognized as a factor of treatment duration. It encompasses missed appointments, compliance with appliance wear, broken appliances, and poor oral hygiene. The role of poor patient cooperation on treatment length has been described in a number of studies.<sup>2,3,5–8</sup> Regardless of patient age, 46% of the variability in treatment duration and 24% of the variability in treatment effectiveness was explained by the number of missed appointments and broken appliances.<sup>6</sup> Poor elastic wear was shown to increase treatment length by a mean of 1.4 months, while three or more poor oral hygiene entries increased treatment time by 1.2 months.<sup>3</sup> Similar findings of the effect of patient cooperation on treatment time have been reported.<sup>9,10</sup>

<sup>1</sup>Private Practice, Las Vegas, Nevada, USA.

<sup>2</sup>Private Practice, Vancouver, British Columbia, Canada.

<sup>3</sup>Associate Clinical Professor, School of Dentistry, University of Nevada, Las Vegas, Nevada, USA.

## CORRESPONDENCE

Dr James Mah  
UNLV School of Dental Medicine  
1001 Shadow Lane  
Las Vegas, NV 89106  
Email: james.mah@sdm.unlv.edu

the PAR weighted scores and the DI when its cephalometric values were deleted. Both the PAR and the DI were determined to be useful indexes for evaluating malocclusion severity and treatment difficulty.<sup>21</sup>

Another major factor influencing both treatment duration and quality is the biomechanical therapy plan. Recently, a novel computer-assisted approach has been introduced.<sup>22</sup> The SureSmile process (OraMetrix) begins with a direct 3D scan of the patient's dentition using an intraoral camera that produces images to create a computer model of the dentition. Various treatment simulations can be performed, and the chosen approach of therapy is used to design and create wires with a bending robot. The deviations of the bends and torques in stainless steel wires are less than 1 degree.<sup>23</sup>

The impact of this system on orthodontic treatment and its duration had yet to be evaluated. Therefore, it was the objective of this study to examine the efficiency and effectiveness of the OraMetrix SureSmile system compared to conventional orthodontic treatment techniques. The hypothesis was that there is a significant difference ( $P < .05$ ) between the OGS scores of the completed SureSmile patients and patients treated by conventional methods.

## MATERIALS AND METHODS

The authors collected the pre- and post-treatment study models of the 62 most recent consecutively completed SureSmile and conventionally treated patients. The pretreatment models were used to determine the DI score and the posttreatment ones to define the OGS score. Treatment length was determined as the time from bonding to debonding. Exclusion criteria were incomplete casts; articulator-mounted casts; or casts from patients with missing molars, orthognathic surgery, or prosthodontic restorations.

Before any scoring, the patients treated with the SureSmile finishing wire ( $n = 38$ ) and those treated traditionally ( $n = 24$ ) were anonymized and randomized. Two independent calibrated examiners (graders) evaluated each model. Statistical analysis

was conducted using the paired sample *t* test, the results of which established the intra- and interexaminer reliability.

The examiners were calibrated using standardized models, the ABO OGS measuring gauge, and the guidelines provided by the ABO OGS.<sup>19</sup> After scoring of each model, the results were reviewed both separately and jointly. Cephalometric values and root parallelism were not appraised as part of the DI in this study. The congruent categories were then scored again on the posttreatment models using the ABO OGS.

## RESULTS

### Data collection analysis

First to be tested using standard diagnostic statistics was whether the data collected by each grader was approximately normally distributed. The distributions had only a slight skew and no kurtosis. This indicated that the data were approximately normally distributed and to be tested with parametric statistical tests.



The mean values for all ABO OGS scores for both examiners are listed in Table 1. Mean value for grader 1 was 27.3 (SD = 7.8) and 28.7 (SD = 8.1) for grader 2. The standard error of the mean (SE), a statistic more appropriate than standard deviation for comparing relatively small samples, was low for the two graders (0.99 and 1.03, respectively) and essentially equal. This suggests that the scores given consistent and therefore reliable.

To compare the measurements of the same patient made by the two graders, the paired samples difference of means test was used. This showed a significant difference ( $P < .05$ ). But a correlation analysis of the two complete sets of measurements showed that the Pearson rank correlation coefficient was 0.96. This suggests that while one grader consistently assigned higher values to a given measurement than the other, the two sets of data were completely consistent with a near one-to-one correspondence. Measurements from both graders showed a similar pattern.

**Table 3** Standard statistics for the individual components of the ABO OGS score of the two treatment modalities

Component	SureSmile		Conventional		Mean difference	Significance (t test, P)
	Mean	SE	Mean	SE		
Alignment and rotations (max)	1.8	0.09	2.7	0.14	0.9	< .05
Alignment and rotations (man)	2.0	0.11	2.3	0.14	0.3	
Marginal ridges (max)	2.0	0.11	2.5	0.17	0.5	< .05
Marginal ridges (man)	2.4	0.12	2.4	0.16	0.0	
Buccolingual inclination (max)	1.2	0.13	1.9	0.18	0.7	< .05
Buccolingual inclination (man)	2.3	0.17	1.9	0.17	-0.4	
Overjet R	2.0	0.15	2.7	0.15	0.7	< .05
Overjet L	2.2	0.14	2.5	0.17	0.3	
Occlusal contacts	3.3	0.18	3.7	0.16	0.4	
Occlusal contacts lingual	3.4	0.18	3.6	0.18	0.2	
Occlusal relationships R	1.8	0.15	2.2	0.18	0.4	
Occlusal relationships L	1.3	0.13	1.3	0.13	0.0	
Interproximal contacts R	0.2	0.05	0.4	0.07	0.2	
Interproximal contacts L	0.2	0.05	0.5	0.09	0.3	< .05

SE = standard error, max = maxillary, man = mandibular.

**Fig 1** Results for the individual components of the ABO OGS score for both treatment modalities and both graders individually.

## DISCUSSION


This study examined the efficiency and effectiveness of the SureSmile system compared to that of conventional fixed appliance straight wire therapy. Two measures were used to evaluate effectiveness: treatment time and ABO OGS score.

The average treatment time for conventionally treated patients in this study was 20.0 months with a standard deviation of 6.40 months. In contrast, in SureSmile patients, it was significantly shorter with an average of 14.7 months. This represents a 36% decrease. The treatment duration for the conventionally

treated patients in this study most closely corresponds with the results reported by Skidmore et al (23.5 months),<sup>3</sup> Fink and Smith (23.1 months),<sup>2</sup> and Alger (22.0 months).<sup>4</sup>

Having identified this difference in treatment length, it was important to find out whether it was due to a different severity of the initial malocclusion. However, the DI scores for both patient groups were similar ( $m = 9.2 \pm 6.6$  for SureSmile,  $m = 11.0 \pm 6.7$  for conventional therapy). Further, the correlation coefficients between ABO OGS and DI were low, suggesting that severity was not a factor impacting these results.

**Fig 2** ABO OGS mean scores for (a) both graders individually and (b) mean score and treatment times (in months) for SureSmile and conventional therapy.



**Table 4** Standard statistics for the treatment time of the two modalities

Modality	Mean	n	SD	SE	Mean difference	SE difference	P
SureSmile	14.7	38	4.71	0.76	-5.3	1.51	<.001
Conventional	20.0	24	6.40	1.21	-5.3	1.51	<.001

n = number of patients, SD = standard deviation, SE = standard error.

**Table 5** Standard statistics for the discrepancy index (DI) of the two modalities

Modality	Mean	n	SD
SureSmile DI	9.2	76	6.58
Conventional DI	11.0	48	6.74

n = number of patients, SD = standard deviation.

Notably, the confidence level for total treatment time with SureSmile is narrower (4.71 SD) than that of conventional treatment (6.40 SD), suggesting that less variation in treatment time is to be expected when treating with SureSmile. This may translate to a better estimation of treatment time.

Part of the standard care process with conventional appliances is that toward the end of therapy, a quality result is generally accomplished through repositioning brackets, altering bracket prescriptions, and/or archwire bending. In contrast, SureSmile prescriptive archwires are derived from an optimal setup. Thus, SureSmile customized archwires overcome the vagrancies of traditional straight-wire appliances and allow for an earlier control. This may be a strong factor for reduced treatment length.

Additionally, SureSmile archwires are bent with high reliability and precision using robotic technology. Unpublished data suggests that the torsional and linear bends are accurate within  $\pm 1$  degree and  $\pm 0.2$  mm, respectively. On the other side, the bracket slots of straight-wire appliances have a very large tolerance, potentially leading to imprecise tooth movements, which are generally corrected by reactive measures resulting in prolonged care.<sup>24,25</sup>

In addition to a shorter care cycle, SureSmile patients demonstrated an OGS score 14.3% better compared to conventional therapy. This finding is important because the prevalent thinking is that better outcomes are related to longer treatment times.<sup>26</sup>

## CONCLUSIONS

The treatment time for the SureSmile system compared to conventional orthodontics was significantly shorter by about 25.0%.

The ABO OGS score for the SureSmile patients was, on average, 14.3% better than for those patients treated with conventional appliances.

## ACKNOWLEDGMENTS

The authors wish to thank Drs Boyd Whitlock, Steve Smith, and Ron Snyder for providing patient records. The authors would also like to thank Harold T. Gross and Michael E. Egan from The Dallas Marketing Group for providing statistical analysis.

## REFERENCES

- Mavreas D, Athanasiou AE. Factors affecting the duration of orthodontic treatment: A systematic review. *Eur J Orthod* 2008;30:386-395.
- Fink DF, Smith RJ. The duration of orthodontic treatment. *Am J Orthod Dentofacial Orthop* 1992;102:45-51.
- Skidmore KJ, Brook KJ, Thomson WM, Harding WJ. Factors influencing treatment time in orthodontic patients. *Am J Orthod Dentofacial Orthop* 2006;129:230-238.
- Alger DW. Appointment frequency versus treatment time. *Am J Orthod Dentofacial Orthop* 1988;94:436-439.
- Beckwith FR, Ackerman RJ Jr, Cobb CM, Tira DE. An evaluation of factors affecting duration of orthodontic treatment. *Am J Orthod Dentofacial Orthop* 1999;115:439-447.
- Robb SI, Sadowsky C, Schneider BJ, BeGole EA. Effectiveness and duration of orthodontic treatment in adults and adolescents. *Am J Orthod Dentofacial Orthop* 1998;114:383-386.
- O'Brien KDO, Robbins R, Vig KWL, Vig PS, Shnorhokian H, Weyant R. The effectiveness of Class II, Division 1 treatment. *Am J Orthod Dentofacial Orthop* 1995;107:329-334.
- Taylor PJ, Kerr WJ, McColl JH. Factors associated with the standard and duration of orthodontic treatment. *Br J Orthod* 1996;23:335-341.
- Shia GJ. Treatment overruns. *J Clin Orthod* 1986;20:602-604.
- Popowich K, Nebbe B, Heo G, Glover KE, Major PW. Predictors for Class II treatment duration. *Am J Orthod Dentofacial Orthop* 2005;127:293-300.
- Cangialosi TJ, Riolo ML, Owens SE Jr, et al. The ABO discrepancy index: A measure of case complexity. *Am J Orthod Dentofacial Orthop* 2004;125:270-278.
- Vu CQ, Roberts WE, Hartsfield JK, Ofner S Jr. Treatment complexity index for assessing the relationship of treatment duration and outcomes in a graduate orthodontics clinic. *Am J Orthod Dentofacial Orthop* 2008;133:9.e1-9.e13.
- Vig KW, Weyant R, Vayda D, O'Brien K, Bennett E. Orthodontic process and outcome: Efficacy studies—strategies for developing process and outcome measures: A new era in orthodontics. *Clin Orthod Res* 1998;1:147-155.
- Richmond S, Shaw WC, Roberts CT, Andrews M. The PAR Index (Peer Assessment Rating): Methods to determine outcome of orthodontic treatment in terms of improvement and standards. *Eur J Orthod* 1992;14:180-187.
- Turbill EA, Richmond S, Wright JL. The time-factor in orthodontics: What influences the duration of treatments in National Health Service practices? *Community Dent Oral Epidemiol* 2001;29:62-72.
- Grewe JM, Hermanson PC. Influence of severity of malocclusion on the duration of orthodontic treatment. *Am J Orthod* 1973;63:533-536.
- Popowich K, Flores-Mir C, Nebbe B, Heo G, Major PW. Comparison of Class I and Class II treatment duration among three different orthodontic practices. *Semin Orthod* 2006;12:52-59.
- Quimby ML, Vig KWL, Rashid RG, Firestone AR. The accuracy and reliability of measurements made on computer-based digital models. *Angle Orthod* 2004;74:298-303.
- Casko JS, Vaden JL, Kokich VG, et al. Objective grading system for dental casts and panoramic radiographs. *Am J Orthod Dentofacial Orthop* 1998;114:589-599.
- Lieber WS, Carlson SK, Baumrind S, Poulton DR. Clinical use of the ABO-Scoring Index: Reliability and subtraction frequency. *Angle Orthod* 2003;73:556-564.
- Deguchi T, Honjo T, Fukunaga T, Miyawaki S, Roberts WE, Takano-Yamamoto T. Clinical assessment of orthodontic outcomes with the peer assessment rating, discrepancy index, objective grading system, and comprehensive clinical assessment. *Am J Orthod Dentofacial Orthop* 2005;127:434-443.
- Mah J, Sachdeva R. Computer-assisted orthodontic treatment: The SureSmile process. *Am J Orthod Dentofacial Orthop* 2001;120:85-87.
- Sachdeva R, Fruge JF, Fruge AM, et al. SureSmile: A report of clinical findings. *J Clin Orthod* 2005;39:297-314.
- Matasa CW. Defend yourself against faulty brackets. *J Gen Orthod* 1991;2:5-9.
- Eliezer R, James LS, Lewis K. Rotation of rectangular wire in rectangular molar tubes: Part I. *Am J Orthod* 1981;80:136-144.
- Sameshima G. Ask us. Some orthodontists claim they can complete treatment relatively quickly, and they rarely treat patients as long as 3 years. How can they do this and still achieve high-quality results. *Am J Orthod Dentofacial Orthop* 2005;127: 526.

suressmile